

Ideal amenability of Banach algebras on locally compact groups

M ESHAGHI GORDJI and S A R HOSSEINIUN*

Department of Mathematics, Faculty of Sciences, Semnan University, Semnan, Iran

*Department of Mathematical Sciences, Shahid Beheshti University, Tehran, Iran

E-mail madjideg@walla.com; r-hosseini@cc.sbu.ac.ir

MS received 25 February 2005; revised 28 June 2005

Abstract. In this paper we study the ideal amenability of Banach algebras. Let $\mathcal{A}$ be a Banach algebra and let I be a closed two-sided ideal in $\mathcal{A}$, $\mathcal{A}$ is I -weakly amenable if $H^1(\mathcal{A}, I^*) = \{0\}$. Further, $\mathcal{A}$ is ideally amenable if $\mathcal{A}$ is I -weakly amenable for every closed two-sided ideal I in $\mathcal{A}$. We know that a continuous homomorphic image of an amenable Banach algebra is again amenable. We show that for ideal amenability the homomorphism property for suitable direct summands is true similar to weak amenability and we apply this result for ideal amenability of Banach algebras on locally compact groups.

Keywords. Amenability; derivation; ideally amenable; weak amenability.

1. Introduction

Let A be a Banach algebra, then A is amenable if $H^1(A, X^*) = \{0\}$ for each Banach A -module X ; this definition was introduced by Johnson in [8]. The Banach algebra A is weakly amenable if $H^1(A, A^*) = \{0\}$; this definition generalizes that introduced by Bade, Curtis and Dales [2]. Let $\mathcal{A}$ be a Banach algebra and let I be a closed two-sided ideal in $\mathcal{A}$, $\mathcal{A}$ is I -weakly amenable if $H^1(\mathcal{A}, I^*) = \{0\}$. $\mathcal{A}$ is n - I -weakly amenable if $H^1(\mathcal{A}, I^{(n)}) = \{0\}$.

Further, $\mathcal{A}$ is ideally amenable if $\mathcal{A}$ is I -weakly amenable for every closed two-sided ideal I in $\mathcal{A}$. This definition was introduced by Eshaghi Gordji and Yazdanpanah in [5]. In this paper we study the ideal amenability of group algebras for a locally compact group.

For example, it was shown in [8] that the group algebra $L^1(G)$ is amenable if and only if G is an amenable group, and in [9] that $L^1(G)$ is weakly amenable for each locally compact group G . We know that there exists a weakly amenable Banach algebra which is not ideally amenable, also every C^* -algebra is ideally amenable [5]. We can not prove that for which locally compact group G , $L^1(G)$ is ideally amenable.

Let A be a Banach algebra. The subspace X of A^* is left introverted if $X^* \cdot X \subseteq X$ and X^* by the following product is a Banach algebra [1]:

$$\langle mn, f \rangle = \langle m, n \cdot f \rangle \quad (m, n \in X^*, f \in X)$$

As in [11] we have the following theorems.

Theorem 1.1. Let A be a Banach algebra and $Y \subseteq X$ two left introverted subspaces of A^* . Then

$$Y^\perp = \{m \in X^*: \langle m, f \rangle = 0 \text{ for every } f \in Y\}$$

is a weak* closed ideal of X^* , also for each $n \in X^*$ and $m \in Y^\perp$ we have $nm = 0$.

In this paper we consider G to be a locally compact (topological) group.

Theorem 1.2. Let X be a left introverted subspace of $L^\infty(G)$ such that $C_0(G) \subseteq X \subseteq CB(G)$. Then there is an isometric algebraic isomorphism $\tau: M(G) \longrightarrow X^*$ and an isometric direct sum decomposition $X^* = \tau(M(G)) \oplus C_0(G)^\perp$ where $C_0(G)^\perp$ is a W^* -closed ideal of X^* and $X^*C_0(G)^\perp = \{0\}$.

Theorem 1.3. Let N be a compact normal subgroup of a locally compact group G , and let $X_N = \{f \in C_0(G): l_h g = f \text{ for } h \in N\}$.

Let $P: C_0(G) \longrightarrow X_N$ be the following linear map:

$$(Pf)(x) = \int_N f(xh)dh,$$

then $P^*: (X_N)^* \longrightarrow M(G)$ is an isometric isomorphism into $M(G)$ and there is the direct sum decomposition $M(G) = P^*(X_N)^* \oplus X_N^\perp$ where $P^*(X_N)^*$ is a closed subalgebra and $X_N^\perp$ is a weak* closed ideal of $M(G)$ contained in right annihilators of $M(G)$.

Let G be a locally compact group and $f \in CB(G)$, then the right orbit of f is given by $R \circ (f) = \{r_x f: x \in G\}$ and let

$$AP(G) = \{f \in CB(G): R \circ (f) \text{ is precompact in the norm topology}\},$$

$$W(G) = \{f \in CB(G): R \circ (f) \text{ is precompact in the weak topology}\}.$$

Theorem 1.4. There is a natural projection $Q: W(G) \longrightarrow AP(G)$ in which $Q^*: AP(G)^* \longrightarrow W(G)^*$ is an isometry and $W(G)^* = Q^*(AP(G)^*) \oplus AP(G)^\perp$ where $Q^*(AP(G)^*)$ is a unital closed subalgebra; and $AP(G)^\perp$ is a weak* closed ideal of $W^*(G)$, and $AP(G)^\perp$ is contained in the right annihilators of $W(G)^*$.

Theorem 1.5. Let G be amenable, and let $A \neq \{0\}$ be a weak* closed self-adjoint translation invariant subalgebra of $L^\infty(G)$. Then there exists a norm one projection $P: L^\infty(G) \longrightarrow A$ which $L^\infty(G)^* = P^*(A^*) \oplus A^\perp$ where $A^\perp$ is a weak* closed ideal of $L^\infty(G)$ and $P^*(A^*)$ is a unital closed subalgebra of $L^\infty(G)^*$ and $A^\perp$ is contained in the right annihilators of $L^\infty(G)^*$.

As in [4] for a left introverted subspace X of PM_p such that $PF_p \subset X \subset UC_p$ we have as follows:

Theorem 1.6. There exists isometric algebra homomorphism $v: W_p \longrightarrow X^*$, such that $X^* = v(W_p) \oplus PF_p^\perp$.

2. Ideal amenability

The concept of ideal amenability was introduced by Eshaghi Gordji and Yazdanpanah [5]. It is well-known that every $\mathcal{C}^*$ -algebra is ideally amenable. A commutative Banach

algebra is ideally amenable if and only if it is weakly amenable. Then ideal amenability is different from amenability. Let $\mathcal{A}$ be a non-unital non-weakly amenable Banach algebra. Then there exists non-inner derivation $D: \mathcal{A} \rightarrow \mathcal{A}^*$. Let $\mathcal{A}^\#$ be the unitization of $\mathcal{A}$, then we define $D_1: \mathcal{A}^\# \rightarrow \mathcal{A}^*$ with $D_1(a + \alpha) = D(a)$, ($a \in \mathcal{A}, \alpha \in \mathbb{C}$). D_1 is a derivation. Since D is not inner, D_1 is not inner. $\mathcal{A}$ is a closed two-sided ideal of $\mathcal{A}^\#$, therefore $\mathcal{A}^\#$ is not ideally amenable. On the other hand, we know that there exists a non-weakly amenable Banach algebra $\mathcal{A}$ such that $\mathcal{A}^\#$ is weakly amenable (see [10]). Thus $\mathcal{A}^\#$ is an example of weakly amenable Banach algebra which is not ideally amenable.

Suppose that $\mathcal{A}$ is a Banach algebra and let X be $\mathcal{A}$ -bimodule. Let $X\mathcal{A} = \{fa: f \in X, a \in \mathcal{A}\}$ and $\mathcal{A}X = \{af: a \in \mathcal{A}, f \in X\}$. The space X is said to be factors on the left (resp. right) if $X = X\mathcal{A}$ (resp. $X = \mathcal{A}X$), and X is neo-unital if X is a factor on both the left and the right. From Theorem 2.3 of [7], we have the following.

Theorem 2.1. *Let $\mathcal{A}$ be a Banach algebra with bounded approximate identity and let I be a codimension 1 closed two-sided ideal of $\mathcal{A}$. Then $H^1(\mathcal{A}, X^*) \cong H^1(I, X^*)$ for every neo-unital Banach $\mathcal{A}$ bimodule X .*

COROLLARY 2.2.

Let G be a discrete group, and let I_0 be a codimension 1 closed two-sided ideal of $l^1(G)$. Then $l^1(G)$ is I_0 -weakly amenable.

Proof. Since $l^1(G)$ is unital then I_0 is neo-unital $l^1(G)$ -bimodule. On the other hand, I_0 is weakly amenable (Prop. 4.2 of [7]). Then by the above theorem $l^1(G)$ is I_0 -weakly amenable.

Let X, Y and Z be normed spaces and let $f: X \times Y \rightarrow Z$ be a continuous bilinear map, then $f^*: Z^* \times X \rightarrow Y^*$ (the transpose of f) is defined by

$$\langle f^*(z^*, x), y \rangle = \langle z^*, f(x, y) \rangle \quad (z^* \in Z^*, x \in X, y \in Y)$$

(f^* is a continuous bilinear map). Clearly, for each $x \in X$, the mapping $z^* \mapsto f^*(z^*, x): Z^* \rightarrow Y^*$ is weak*-weak* continuous. Let $f^{**} = (f^*)^*$ and $f^{***} = (f^{**})^*$, then $f^{***}: X^{**} \times Y^{**} \rightarrow Z^{**}$ is the unique extension of f such that $f^{***}(\cdot, y^{**}): X^{**} \rightarrow Z^{**}$ is weak*-weak* continuous. Also $f^{***}(x, \cdot): Y^{**} \rightarrow Z^{**}$ is weak*-weak* continuous for every $x \in X$. Let $f^r: Y \times X \rightarrow Z$ be defined by $f^r(y, x) = f(x, y)$, ($x \in X, y \in Y$). Then f^r is a continuous linear mapping from $Y \times X$ to Z . The map f is Arens regular whenever $f^{***} = f^{r***r}$. This is equivalent to the condition that the map $f^{***}(x^{**}, \cdot): Y^{**} \rightarrow Z^{**}$ be weak*-weak* continuous for every $x^{**} \in X^{**}$. Let I be a closed two-sided ideal of $\mathcal{A}$ and $\pi_l: \mathcal{A} \times I \rightarrow I$ and $\pi_r: I \times \mathcal{A} \rightarrow I$ are the right and left module actions of $\mathcal{A}$ on I respectively. Then I^{**} is a $\mathcal{A}^{**}$ -bimodule with module actions $\pi_l^{***}: \mathcal{A}^{**} \times I^{**} \rightarrow I^{**}$ and $\pi_r^{***}: I^{**} \times \mathcal{A}^{**} \rightarrow I^{**}$. π_l is Arens regular if and only if a'' is fixed in $\mathcal{A}^{**}$. The mapping $i'' \mapsto \pi_l^{***}(i'', a''): I^{**} \rightarrow I^{**}$ is weak*-weak* continuous and π_r is Arens regular if and only if i'' is fixed in I^{**} . The mapping $a'' \mapsto \pi_r^{***}(a'', i''): \mathcal{A}^{**} \rightarrow I^{**}$ is weak*-weak* continuous. We now prove the following lemma.

Lemma 2.3. *Let $\mathcal{A}$ be a Banach algebra with bounded approximate identity and let I be a closed two-sided ideal of $\mathcal{A}$. The the following assertions hold.*

- (i) *If $\pi_r: I \times \mathcal{A} \rightarrow I$ be Arens regular, then I^* factors on the right.*
- (ii) *If $\pi_l: \mathcal{A} \times I \rightarrow I$ be Arens regular, then I^* factors on the left.*

Proof. Let (e_α) be a bounded approximate identity for $\mathcal{A}$ with cluster point E . Suppose that π_r be Arens regular and let $i'' \in I^{**}$ be the cluster point of (i_β) ((i_β) is a net in I). Then we have

$$\begin{aligned} i'' &= \lim_{\beta} \lim_{\alpha} i_{\beta} \\ &= \lim_{\beta} \lim_{\alpha} i_{\beta} e_{\alpha} \\ &= \lim_{\alpha} \lim_{\beta} i_{\beta} e_{\alpha} \\ &= \lim_{\alpha} i'' e_{\alpha} = i'' E. \end{aligned}$$

Then for $i' \in I^*$ we have

$$\begin{aligned} \lim_{\alpha} \langle e_{\alpha} i', i'' \rangle &= \lim_{\alpha} \langle i', i'' e_{\alpha} \rangle \\ &= \langle i', i'' E \rangle \\ &= \langle i', i'' \rangle. \end{aligned}$$

Thus $e_{\alpha} i' \rightarrow i'$ weakly. Since $e_{\alpha} i' \in \mathcal{A} I^*$ for every α , by the Cohen–Hewit factorization theorem we know that $\mathcal{A} I^*$ is closed in I^* , then $i' \in \mathcal{A} I^*$. Thus the proof of (i) is complete. For (ii), let $i'' \in I^{**}$ be the cluster point of (i_{β}) . For each β we know that $E i_{\beta} = i_{\beta}$ since π_l is Arens regular, then $e_{\alpha} i'' \rightarrow E i'' = i''$ by weak* topology of I^{**} . Then for every $i' \in I^*$,

$$\begin{aligned} \lim_{\alpha} \langle i' e_{\alpha}, i'' \rangle &= \lim_{\alpha} \langle i', e_{\alpha} i'' \rangle \\ &= \langle i', E i'' \rangle \\ &= \langle i', i'' \rangle. \end{aligned}$$

Therefore $i' e_{\alpha} \rightarrow i'$ weakly. Again by the Cohen–Hewit factorization theorem we conclude that I^* factors on the left.

COROLLARY 2.4.

Let $\mathcal{A}$ be a Banach algebra with bounded approximate identity and let I be a codimension 1 closed two-sided ideal of $\mathcal{A}$. If the module actions of $\mathcal{A}$ on I are Arens regular, then $\mathcal{A}$ is 2- I -weakly amenable if and only if I is 2-weakly amenable.

3. Ideal amenability and direct sum decompositions

We prove an elementary lemma about derivations and homomorphic images and apply this to ideal amenability for various algebras defined over locally compact groups.

Lemma 3.1. *Let A be a right unital Banach algebra and $A = B \oplus I$ for closed unital sub-algebra B and closed two-sided ideal I in A where I is contained in the set of right annihilators of A . Then we have the following:*

- (i) *for every closed two-sided ideal J in B , $J \oplus I$ is a closed two-sided ideal in A ,*
- (ii) *if A is ideally amenable, then B is ideally amenable.*

Proof.

- (i) First we show that $J \oplus I$ is a two-sided ideal in A . Let $a = b + i \in A$ and $x = j + i' \in J \oplus I$ since $AI = \{0\}$, then we have

$$Ax = (b + i)(j + i') = bj + ij \in J \oplus I,$$

$$xa = (j + i')(b + i) = jb + i'b \in J \oplus I.$$

Let e be a right unit element in A , then there exists $b_1 \in B$ and $i_1 \in I$ such that $e = b_1 + i_1$. Then $e = e^2 = e(b_1 + i_1) = eb_1$ and b_1 is a right unit element for A . But B is unital, and so $b_1 = 1_B$. Let $\|1_B\| = M$ and $j + i \in J + I$, then we have

$$\|j + i\| \geq M^{-1}\|1_B(j + i)\| = M^{-1}\|j\|.$$

Then the natural map $J \longrightarrow \frac{J+I}{I}$ is bicontinuous, therefore $(J + I)/I$ is complete. J is closed in A/I , and $J + I$ is closed in A .

- (ii) Let J be a closed ideal in B and $D: B \longrightarrow J^*$ be a derivation and $\pi: B \oplus I \longrightarrow B$ be the natural embedding. π is a homomorphism and so is $\pi|_{J \oplus I}$. We show that $\bar{D} = (\pi|_{J \oplus I})^* \circ D \circ \pi: A \longrightarrow (J \oplus I)^*$ is a derivation. For $x, y \in A$ and $z \in J \oplus I$, we have

$$\begin{aligned} \langle (\pi|_{J \oplus I})^* \circ D \circ \pi(xy), z \rangle &= \langle D(\pi(x)\pi(y)), \pi(z) \rangle \\ &= \langle D(\pi(x)) \cdot \pi(y) + \pi(x) \cdot D(\pi(y)), \pi(z) \rangle, \end{aligned}$$

but

$$\begin{aligned} \langle D(\pi(x)) \cdot \pi(y), \pi(z) \rangle &= \langle D(\pi(x)), \pi(y)\pi(z) \rangle \\ &= \langle D(\pi(x)), \pi(yz) \rangle \\ &= \langle \bar{D}(x), yz \rangle \\ &= \langle \bar{D}(x) \cdot y, z \rangle. \end{aligned}$$

Similarly,

$$\langle \pi(x) \cdot D(\pi(y)), \pi(z) \rangle = \langle x \cdot \bar{D}(y), z \rangle,$$

then $\bar{D}(xy) = x \cdot \bar{D}(y) + \bar{D}(x) \cdot y$ and $\bar{D}$ is a derivation. A is ideally amenable and by (i), $J \oplus I$ is a closed two-sided ideal in A . Then there exists $\xi \in (J \oplus I)^*$ such that $\bar{D} = \delta_\xi$, and let $\eta = \xi|_J$, then for $b \in B$ and $j \in J$ we have

$$\begin{aligned} \langle Db, j \rangle &= \langle D(\pi(b)), \pi(j) \rangle \\ &= \langle \bar{D}b, j \rangle \\ &= \langle b\xi - \xi \cdot b, j \rangle \\ &= \langle \xi, j \cdot b \rangle - \langle \xi, b \cdot j \rangle \\ &= \langle b \cdot \eta - \eta \cdot b, j \rangle \\ &= \delta_\eta(b)(j). \end{aligned}$$

Therefore $D = \delta_\eta$ and B is ideally amenable.

COROLLARY 3.2.

Let $\mathcal{A}$ be a dual Banach algebra with bounded approximate identity and let $\mathcal{A}^{**}$ be the second dual of $\mathcal{A}$ with the first Arens product. If $\mathcal{A}^{**}$ be ideally amenable then $\mathcal{A}$ is ideally amenable.

Proof. Let $\mathcal{A}_*$ be the pre-dual of $\mathcal{A}$. Then it is easy to see that $\mathcal{A}^{**} = \mathcal{A} \oplus (\mathcal{A}_*)^\perp$ (see, for example, [3] or [6]). On the other hand $\mathcal{A}$ has bounded approximate identity, then $\mathcal{A}^{**}$ has right identity. By lemma the ideal amenability of $\mathcal{A}^{**}$ implies that $\mathcal{A}$ is also ideally amenable.

We know that $M(G)$ is a dual Banach algebra (with pre-dual $C_0(G)$). Thus by Corollary 3.2, $M(G)$ is ideally amenable whenever $M(G)^{**}$ is ideally amenable.

COROLLARY 3.3.

Let X be a left introverted subspace of $L^\infty(G)$, such that $C_0(G) \subseteq X \subseteq CB(G)$, if X^* be ideally amenable, then $M(G)$ and $L^1(G)$ are ideally amenable.

Proof. By applying Lemma 2.1 and Theorem 1.2, ideal amenability of X^* implies that $M(G)$ is ideally amenable, and by [5] we know that if $M(G)$ is ideally amenable, then $L^1(G)$ is also ideally amenable.

COROLLARY 3.4.

Let N be a compact normal subgroup of a locally compact group G , and X_N and P are as in Theorem 1.3. Then ideal amenability of $M(G)$ implies that $P^*(X_N)^*$ is ideally amenable.

COROLLARY 3.5.

In Theorem 1.4, if $W(G)^*$ be ideally amenable, then $Q^*(AP(G)^*)$ is ideally amenable.

COROLLARY 3.6.

In Theorem 1.5, if $L^\infty(G)^*$ be ideally amenable, then $P^*(A^*)$ is ideally amenable.

Let E be a right identity of $L^1(G)^{**}$ with $E \geq 0$ and $\|E\| = 1$, then we have the isomorphism $EL^1(G)^{**} \simeq LUC(G)^*$ and we have the decomposition

$$\begin{aligned} L^1(G)^{**} &= EL^1(G)^{**} \oplus (I - E)L^1(G)^{**} \\ &= LUC(G)^* \oplus (I - E)L^1(G)^{**}, \end{aligned}$$

where $(I - E)L^1(G)^{**}$ is contained in the right annihilators of $L^1(G)^{**}$. So we have the decomposition $LUC(G)^* = M(G) \oplus C_0(G)^\perp$ where $C_0(G)^\perp$ is a closed ideal in $LUC(G)^*$ and contained in the right annihilators of $LUC(G)^*$ (see, for example, [11]). Therefore the following proposition follows easily from Theorem 2.1.

PROPOSITION 3.7.

- (i) Suppose that $L^1(G)^{**}$ is ideally amenable, then $LUC(G)^*$ is ideally amenable.
- (ii) Suppose $LUC(G)^*$ is ideally amenable, then $M(G)$ is ideally amenable.
- (iii) Suppose $M(G)$ is ideally amenable, then $L^1(G)$ is ideally amenable.

By Theorem 1.6 we have as follows.

PROPOSITION 3.8.

Let X be a left introverted subspace of PM_p , such that $PM_p \subseteq X \subseteq UC_p$, if X^* be ideally amenable, then W_p is ideally amenable.

References

- [1] Arens R, The adjoint of a bilinear operation, *Proc. Am. Math. Soc.* (2) (1951) 839–848
- [2] Bade W G, Curtis P G and Dales H G, Amenability and weak amenability for Beurling and Lipschitz algebras, *Proc. London Math. Soc.* **55**(3) (1987) 359–377
- [3] Dales H G and Lau A T M, The second duals of Beurling algebras, preprint
- [4] Derighetti A, Filali M and Sangani Monfared M, On the ideal structure of some Banach algebras related to convolution operators on $L^p(G)$, to appear
- [5] Eshaghi Gordji M and Yazdanpanah T, Derivations into duals of ideals of Banach algebras, *Proc. Ind. Acad. Sci. (Math. Sci.)* **114**(4) (2004) 399–403
- [6] Ghahramani F and Laali J, Amenability and topological centers of the second duals of Banach algebras, *Bull. Austral. Math. Soc.* **65** (2002) 191–197
- [7] Gronbaek N and Lau A T, On hochschild cohomology of the augmentation ideal of a locally compact group, *Math. Proc. Comb. Soc.* **126** (1999) 139–148
- [8] Johnson B E, Cohomology in Banach algebras, *Mem. Am. Math. Soc.* **127** (1972)
- [9] Johnson B E, Weak amenability of group algebras, *Bull. London Math. Soc.* **23** (1991) 281–284
- [10] Johnson B E and White M C, A non-weakly amenable augmentation ideal, to appear
- [11] Lau A T and Loy R J, Weak amenability of Banach algebras on locally compact groups, *J. Funct. Anal.* **145**(1) (1997) 175–204